arXiv:math/0405022v1 [math.GR] 2 May 2004

THE TITS ALTERNATIVE FOR CAT(0) CUBICAL COMPLEXES

MICHAH SAGEEV AND DANIEL T. WISE

Abstract. We prove a Tits alternative theorem for groups acting on CAT(0) cubical complexes. Namely, suppose that G is a group for which there is a bound on the orders of its finite subgroups. We prove that if G acts properly on a finite-dimensional CAT(0) cubical complex, then either G contains a free subgroup of rank 2 or G is finitely generated and virtually abelian. In particular the above conclusion holds for any group G with a free action on a finite-dimensional CAT(0) cubical complex.

1. Introduction

A group G satisfies the *Tits alternative* if for every subgroup H of G, either H is virtually solvable or H contains a free subgroup of rank 2. The Tits alternative is named for Jacque Tits, who discovered that it holds for any finitely generated linear group [Tit72]. The Tits alternative has many important consequences - for instance, it implies von Neumann's dichotomy between amenability and containing a free subgroup of rank 2.

The purpose of this note is to prove:

Theorem 1.1. *Suppose that G is a group for which there is a bound on the order of its finite subgroups. Suppose that G acts properly on a CAT(0) cubical complex. Then for each subgroup H of G, either H contains a rank 2 free subgroup, or H is virtually a finitely generated abelian group.*

Since finite subgroups of G have fixed points in any action of G on a CAT(0) space, we have the following corollary.

Corollary 1.2. *Let G be a group which acts freely on a finite dimensional CAT(0) cubical complex. Then for each subgroup H of G, either H contains a rank 2 free subgroup, or H is virtually a finitely generated abelian group.*

Theorem 1.1 is false without the finite dimensional hypothesis. Indeed, Thompson's group is not virtually solvable and contains no rank 2 free subgroup [BS85], but Thompson's group acts freely on an infinite dimensional (yet proper) CAT(0) cubical complex [Far03].

Theorem 1.1 also fails to hold without a bound on the order of finite subgroups. Indeed, let G be the ascending union of finite groups $G = G_1 \subset G_2 \subset \cdots$ (e.g. the direct product of countably many finite groups). Then there exists a "coset tree"

Date: February 1, 2008.

2000 *Mathematics Subject Classification.* 20F67, 20E08 .

Key words and phrases. CAT(0) cubical complex, Tits alternative.

Sageev's research supported by BSF and ISF.

Wise's research supported by FCAR and NSERC.

T on which G acts (see [Ser80]); namely, the lattice of left cosets of the subgroups G_i is a tree. The natural left action of G on T is proper, but G does not satisfy the conclusion of Theorem 1.1.

We list some cases in which the Tits alternative is already known for groups arising in geometric group theory. Note that G satisfies the *weak Tits alternative* if either G is virtually solvable, or G contains F_2 (as opposed to requiring this for each finitely generated subgroup).

The Tits Alternative holds for the following groups:

1. word-hyperbolic groups [Gro87]
2. $\mathrm{Out}(F_n)$ [BFH00]
3. Foldable cubical chamber complexes [BŚ99]
4. $\pi_1 M$ where M is a nonpositively curved real-analytic 4-manifold satisfying a certain cycle condition [Xie04]
5. fundamental groups of CAT(0) square complexes with no fake planes [Xie]

The Weak Tits Alternative holds for π_1 of a compact space satisfying:

6. $C(4)$-$T(4)$: [Col73]
7. $C(6)$: [EM86] (further discusses the $C(4)$-$T(4)$ case and references $C(6)$ case to dissertation in bibliography)
8. $C(3)$-$T(6)$: [EH88]
9. Finite 2-complexes satisfying the Gersten-Pride weight test with positive rational angles [CT96]
10. Finite nonpositively curved 2-complexes [BB95]

The results most closely related to this paper are those of Ballman and Świątkowski [BŚ99] and Xie [Xie]. We note that Ballmann and Świątkowski actually proved a stronger result allowing larger stabilizers in a more restrictive setting. We also note that Theorem 1.1 unifies some of the results described above in the small-cancellation case. Indeed, fundamental groups of compact $C'(\frac{1}{6})$ and $C'(\frac{1}{4})$-$T(4)$ complexes act with uniformly bounded stabilizers on finite dimensional CAT(0) cube complexes [Wis]. These are the most important metric subcases of the $C(6)$ and $C(4)$-$T(4)$ groups above.

We close the introduction with the following problems:

Problem 1.3. Does the Tits alternative hold for G if:

1. G acts properly discontinuously and cocompactly on a CAT(0) space? (This is even unknown for CAT(0) manifolds.)
2. G is automatic?

2. Preliminaries

2.1. **Cubical complexes and hyperplanes.** We recall some basic facts about hyperplanes (see [Sag95] for details). Let X be a CAT(0) cubical complex. Two edges e and f are *square equivalent* if they are opposite edges of a square σ in X. We let $\sim$ denote the equivalence relation generated by square equivalence and let $\overline{e}$ denote the equivalence class containing e. We identify each n-cube σ of X with the standard unit cube in $\mathbb{R}^n$. A *midcube* of σ is the intersection of σ with an $(n-1)$-dimensional hyperplane parallel to one of the faces of σ and containing the center of σ. A *hyperplane* in X is the union of all midcubes meeting a particular equivalence class of edges. We note some basic facts about hyperplanes.

1. A hyperplane meets each cube in at most one midcube.
2. A hyperplane separates X into precisely two components.
3. Each hyperplane is itself a cubical complex, and inherits a CAT(0) structure from X.

Let H be a subgroup of the finitely generated group G. The number $e(G, H)$ of *ends of G with respect to H* is defined to be the number of ends of the coset graph of G relative to H.

In [Sag95], the following theorem is proven

Theorem 2.1. *Suppose that G acts on a finite dimensional $CAT(0)$ cubical complex without a global fixed point. Then there exists a hyperplane J in X such that $e(G, stab(J)) > 1$.*

This result was later generalized by Niblo and Roller [NR98] and Gerasimov [Ger97] to actions on infinite dimensional complexes.

2.2. **The Algebraic Torus Theorem.** We will appeal to the following theorem of Dunwoody and Swenson [DS00], which is a generalization of a theorem of Scott and Swarup [SS00] and Bowditch [Bow98]. The following statement is a slightly weakened version of the statement appearing in [DS00].

Theorem 2.2 (The Algebraic Torus Theorem)**.** *Suppose that G is a finitely generated group which contains a virtually polycyclic subgroup H with $e(G, H) > 1$. Then one of the following holds:*

1. *G is virtually polycyclic.*
2. *G has a non-elementary fuchsian quotient with virtually polycyclic kernel.*
3. *G splits as a free product with amalgamation or HNN-extension over a virtually polycyclic subgroup.*

2.3. **Groups acting on CAT(0) spaces.** Finally, we record the following important facts about groups acting on CAT(0) spaces.

Lemma 2.3. *(1) If G acts cellularly on a CAT(0) complex X with finitely many shapes, then G acts semi-simply with a discrete set of translation lengths* [Bri99]. *(2) If G acts semi-simply and properly-discontinuously on the CAT(0) space X, then any virtually solvable subgroup H of G is virtually abelian* [BH99].

Moreover, we have the following theorem (see [BH99, Thm 7.5 and Rem 7.7]):

Theorem 2.4. *Let G act by semi-simple isometries on a CAT(0) space so that:*

1. *there is a bound on the dimension of an isometrically embedded flat,*
2. *the set of translation numbers of elements of G is discrete at 0,*
3. *there is a bound on the order of finite subgroups.*

Then any sequence $H_1 \subsetneq H_2 \subsetneq \cdots$ of virtually abelain subgroups terminates.

Corollary 2.5. *Let G act as in Theorem 2.4. If $H^t \subset H$ then $H^t = H$.*

Proof. This follows from Theorem 2.4 by setting $H_i = H^{t^{-i}}$. □

3. The Main Theorem

We now prove Theorem 1.1 in the following form:

Theorem 3.1. *Suppose there is a uniform bound on the orders of finite subgroups of the group G. Let G act properly on a finite dimensional CAT(0) cubical complex X. Then either G has a rank 2 free subgroup or G is virtually a finitely generated abelian group.*

Proof. We proceed by induction on $\dim(X)$. If $\dim(X) = 0$ then G is finite and hence virtually finitely generated abelian. Suppose the theorem holds for $\dim(X) < n$, and consider the case where $\dim(X) = n$.

We first prove the theorem in the case that G is finitely generated. If G is finite, we are done, so suppose that G is infinite. Since G is infinite and acts properly on X, it has no global fixed point. Therefore, by Theorem 2.1, there exists a hyperplane J in X with $e(G, stab(J)) > 1$. Let $H = stab(J)$. Then H acts properly on J. By induction, either H is virtually finitely generated abelian or H contains a rank 2 free subgroup. If H has a rank 2 free subgroup then so does G and we are done. We therefore assume that H is virtually a finitely generated abelian group. Applying Theorem 2.2 we have either:

1. G is virtually polycyclic,
2. G has a non-elementary fuchsian quotient with virtually polycyclic kernel,
3. G splits as a free product with amalgamation or HNN-extension over a virtually polycyclic subgroup.

In case (1), G is virtually abelian by Lemma 2.3(2). In case (2), the non-elementary fuchsian quotient contains a rank 2 free subgroup, and hence G does as well. Thus, suppose that G splits as an amalgamated free product $A *_P B$ or HNN extension $C*_P$ over a virtually polycyclic group P. Since X has finitely many shapes, G acts semi-simply by Lemma 2.3(1), and so P is virtually abelian by Lemma 2.3(2).

An application of the normal form theorem for graphs of groups [Ser80], shows that if either $[A : P] > 2$ or $[B : P] > 2$ then G contains a rank 2 free subgroup. We may therefore assume that $[A : P] = 2$ and $[B : P] = 2$. In this case G is virtually polycyclic. Indeed, the Bass-Serre tree is a line, so G has an index 2 subgroup G' which acts by translations, and the resulting homomorphism $G' \to \mathbb{Z}$ has kernel P, so $G' \cong P \ltimes \mathbb{Z}$.

Again, the normal form theorem shows that if both $[C : P] > 1$ and $[C : P^t] > 1$ (where t is the stable letter of $C*_P$) then G contains a rank 2 free subgroup. On the other hand, by Corollary 2.5, if $[C : P] = 1$ then $[C : P^t] = 1$ (and vice-versa). We may therefore assume that $[C : P] = 1$ and $[C : P^t] = 1$, and so $G \cong P \ltimes \mathbb{Z}$.

In each case, G is virtually polycyclic and hence virtually abelian by Lemma 2.3.

Now suppose that G is not finitely generated. Thus, G contains an infinite ascending sequence of proper subgroups $G_1 \subsetneq G_2 \subsetneq G_3 \dots$. We are done if any G_i contains a rank 2 free subgroup. So we shall show that the assumption that each G_i is virtually finitely generated abelian leads to a contradiction. We first verify conditions (1)-(3) in the hypothesis of Theorem 2.4. Condition (1) holds since X is finite dimensional. Condition (2) holds by Lemma 2.3, since X is a finite dimensional cubical complex and hence a polyhedral complex with finitely many shapes. Condition (3) is hypothesized in our theorem. Thus the infinite sequence $G_1 \subsetneq G_2 \subsetneq G_3 \ldots$ terminates by Theorem 2.4 which is impossible. $\square$

References

[BB95] Werner Ballmann and Michael Brin. Orbihedra of nonpositive curvature. *Inst. Hautes Études Sci. Publ. Math.*, (82):169–209 (1996), 1995.

[BFH00] Mladen Bestvina, Mark Feighn, and Michael Handel. The Tits alternative for $\mathrm{Out}(F_n)$. I. Dynamics of exponentially-growing automorphisms. *Ann. of Math. (2)*, 151(2):517–623, 2000.

[BH99] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature.* Springer-Verlag, Berlin, 1999.

[Bow98] Brian H. Bowditch. Cut points and canonical splittings of hyperbolic groups. *Acta Math.*, 180(2):145–186, 1998.

[Bri99] Martin R. Bridson. On the semisimplicity of polyhedral isometries. *Proc. Amer. Math. Soc.*, 127(7):2143–2146, 1999.

[BS85] Matthew G. Brin and Craig C. Squier. Groups of piecewise linear homeomorphisms of the real line. *Invent. Math.*, 79(3):485–498, 1985.

[BŚ99] Werner Ballmann and Jacek Świątkowski. On groups acting on nonpositively curved cubical complexes. *Enseign. Math. (2)*, 45(1-2):51–81, 1999.

[Col73] Donald J. Collins. Free subgroups of small cancellation groups. *Proc. London Math. Soc. (3)*, 26:193–206, 1973.

[CT96] J. M. Corson and B. Trace. Geometry and algebra of nonspherical 2-complexes. *J. London Math. Soc. (2)*, 54(1):180–198, 1996.

[DS00] M. J. Dunwoody and E. L. Swenson. The algebraic torus theorem. *Invent. Math.*, 140(3):605–637, 2000.

[EH88] Martin Edjvet and James Howie. Star graphs, projective planes and free subgroups in small cancellation groups. *Proc. London Math. Soc. (3)*, 57(2):301–328, 1988.

[EM86] Mohamed S. El-Mosalamy. Free subgroups of small cancellation groups. *Israel J. Math.*, 56(3):345–348, 1986.

[Far03] Daniel S. Farley. Finiteness and CAT(0) properties of diagram groups. *Topology*, 42(5):1065–1082, 2003.

[Ger97] V. N. Gerasimov. Semi-splittings of groups and actions on cubings. In *Algebra, geometry, analysis and mathematical physics (Russian) (Novosibirsk, 1996)*, pages 91–109, 190. Izdat. Ross. Akad. Nauk Sib. Otd. Inst. Mat., Novosibirsk, 1997.

[Gro87] M. Gromov. Hyperbolic groups. In *Essays in group theory*, volume 8 of *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New York, 1987.

[NR98] Graham A. Niblo and Martin A. Roller. Groups acting on cubes and Kazhdan's property (T). *Proc. Amer. Math. Soc.*, 126(3):693–699, 1998.

[Sag95] Michah Sageev. Ends of group pairs and non-positively curved cube complexes. *Proc. London Math. Soc. (3)*, 71(3):585–617, 1995.

[Ser80] Jean-Pierre Serre. *Trees.* Springer-Verlag, Berlin, 1980. Translated from the French by John Stillwell.

[SS00] G. P. Scott and G. A. Swarup. An algebraic annulus theorem. *Pacific J. Math.*, 196(2):461–506, 2000.

[Tit72] J. Tits. Free subgroups in linear groups. *J. Algebra*, 20:250–270, 1972.

[Wis] Daniel T. Wise. Cubulating small-cancellation groups. *GAFA*. To appear.

[Xie] Xiangdong Xie. Groups acting on CAT(0) square complexes. Preprint 2003.

[Xie04] Xiangdong Xie. Tits alternative for closed real analytic 4-manifolds of nonpositive curvature. *Topology Appl.*, 136(1-3):87–121, 2004.

Dept. of Math., Technion, Haifa 32000, Israel
E-mail address: `sageevm@techunix.technion.ac.il`

Math. & Stats., McGill University, Montreal, Quebec, Canada H3A 2K6
E-mail address: `wise@math.mcgill.ca`